

ATSR

Algorithm, Cloud cover, height and temperature

ORAC Oxford/RAL retrieval of aerosol and clouds

Caroline Poulsen, Andy Sayer³, Richard Siddans,
Barry Latter and, Chris Arnold², Don Grainger²(PI),
Gareth Thomas² and Phil Watts¹

Space Science and Technology Department
Rutherford Appleton Laboratory
UK

1. *Now at EUMETSAT*
2. *University of Oxford*
3. *UO/RAL*







Science & Technology Facilities Council
Rutherford Appleton Laboratory




Time line of ATSR (vis/nir and IR) instruments

1985 1990 1995 2000 2005 2010 2015





ATSR Scan Geometry and Channels

- Dual-viewing visible & IR radiometers
 - Nadir 0° - 22° , forward 53° - 56° , 100 seconds difference
 - 550 nm, 660 nm, 870 nm, $1.6\text{ }\mu\text{m}$, $3.7\text{ }\mu\text{m}$, $10.8\text{ }\mu\text{m}$, $12\text{ }\mu\text{m}$ (1.38,2.25)



The ATSR scan geometry (courtesy ESA).

ATSR-2/AATSR/SLSTR



ATSR-2

1km resolution

512km swath over land

300km swath over sea

LTDN 10:30

AATSR

1km resolution

512km swath over land

512km swath over sea

LTDN 10:00

SLSTR

0.5-1km resolution

Nadir swath 1300 - 1800 km

Dual view swath 750 km

LTDN 10:00

SOME NOMENCLATURE

- **ORAC:** Oxford – Rutherford Appleton Laboratory (RAL) Aerosol & Cloud retrieval scheme
- **GRAPE:** Global Retrieval of ATSR Cloud Parameters & Evaluation
 - Project in which ORAC first applied to generate a global cloud dataset, 1995 onwards
 - Cloud-free scenes processed with simple aerosol retrieval
 - Available from <http://badc.nerc.ac.uk/data/grape/>

OPTIMAL ESTIMATION: BASICS


- Find most probable state $\mathbf{x}$ given:
 - Measurements $\mathbf{y}$ with covariance $\mathbf{S}_y$
 - Any *a priori* information $\mathbf{x}_a$ with covariance $\mathbf{S}_a$
- Simultaneous fit of all cloud properties to observations in all spectral channels
 - Physically consistent between different spectral domains
- Robust error propagation
 - Measurement, forward model, *a priori*
 - Goodness-of-fit ('cost') statistic on each retrieval
 - Uncertainty estimates $\mathbf{S}_x$ on retrieved state variables
- Can extend easily to different sensors or combinations of sensors, provided uncertainties can be characterised

ORAC CLOUDS: OVERVIEW

- Algorithm: Poulsen *et al.*, (in prep 2010, P. Watts *et al* 1997 EUMETSAT report)
- Validation: Sayer *et al.* (in prep)
- Separate solar/thermal forward models
 - 660 nm, 870 nm, 1.6 μm
 - 10.8 μm , 12 μm
 - Consistent ice (Baran) & water properties propagated through
- Single cloud layer
- Simple split between cloudy/clear portions


Retrieved in state vector	Derived
Cloud optical depth at 550 nm	Liquid/ice water path
Cloud effective radius	
(Phase)	
Cloud-top pressure	Cloud-top height
	Cloud-top temperature
Surface temperature	

SOLAR FORWARD MODEL: CLOUDS



- Lookup tables (LUTs) of atmospheric transmittance, reflectance (T, R)
 - DISORT radiative transfer model
 - MODTRAN gas absorption
- Lambertian surface albedo R_s
 - MODIS albedo product for 2002 over land
 - Cox & Munk (1954) glint over sea, ECMWF 10 m winds
- Rayleigh scattering/gas absorption above & below cloud layer
 - ECMWF $p/T/H_2O$ profiles


THERMAL FORWARD MODEL: CLOUDS



$$R = R_{ac}^{\uparrow} + R_{ac}^{\downarrow} R_D T_{ac} + R_{bc}^{\uparrow} T_D T_{ac} + B(T(p_c)) E_m T_{ac}$$

- Similar in principle to solar forward model
 - ECMWF $p/T/H_2O$ profiles & surface temperature
 - RTTOV radiative transfer model


EXAMPLE SCENE



EXAMPLE SCENE

Images from Poulsen *et al.*, 2010







- Cost indicator of where model not consistent with observations
- Errors on cloud-top pressure increase as clouds get thinner.




Cost

Cloud top pressure

Uncertainty




EXAMPLE SCENE





- Phase corresponds to what expected from false-colour image
- Cloud fraction largely determined by ATSR cloud mask
 - On 3x4 km scale, most cloudy scenes fully overcast

Multi-layer Simulations



Non Linear retrieval simulations of **multi layer** clouds using the ORAC cloud model.


Low cloud is at 780hPa and high cloud is at 320hPa, optical depth of cloud layers is varied.




Non Linear retrieval
simulations of **multi layer**
clouds using the ORAC
cloud model.

Quality control applied to
GEWEX files is cost <10

Identifies many multi-layer
cloud conditions but
not all – some heights still
radiative average of 2
layers




Example Retrieval of multi layer cloud





Example of OCA algorithm applied to SEVIRI data and compared with collocated Calipso and Cloud sat data

- IR channels have a penetration depth of approx 1 optical depth
- CTP has a negative bias.

Global location of multi layered cloud






High values indicate the presence of multi layered cloud




Relative cloud amounts: Comparison with other satellites

— ATSR2
— AATSR
— PATMOSX
— ISCCP
— AIRS
— MODIS-STT
— MODIS-CET
— MISR
— POLDER
— CALIPSO

NB. Satellites are collocated with ATSR




Temperature inversion problem



Cloud fraction: Comparison with other satellites


Total cloud fraction: SON




- ATSR2
- AATSR
- PATMOSX
- ISCCP
- AIRS
- MODIS-STT
- MODIS-CET
- MISR
- POLDER
- CALIPSO

**NB. Satellites
are
collocated
with ATSR**



CALIPSO-GOCCP




ATSR-2





Comparison of ATSR cloud top pressure with other satellite sensors





Multi annual mean CTP comparison



- Problems with the boundary layer inversions
- Underestimation of altitude in the tropics due to IR penetration
- Problems with multi layer cloud
- Sampling is different




**AATSR detects
thicker high
cloud**



**AATSR misses
thin cloud**

Cloud top temperature




ATSR CTT 3.6K cooler than MODIS-CE
ATSR CTT 6.4K cooler than MODIS-ST

Spread in CTT is smaller than in CTP (about 1km compared to 3km in CTP)

- **Although CTH may be too high, e.g temperature inversions, CTT may be correct.**
- **Sampling likely to be a large contributing factor to differences**

Conclusions


- ATSR detects cloud well over sea, cloud detection over land could be improved.
- Negative bias in cloud top height is expected when observing multi layered cloud, this cloud is often identified using the cost threshold.
 - ATSR detects too much mid level cloud
 - Problem with temperature inversion
 - To be corrected in next processing
 - Could apply quick fix to existing data
- Differences in cloud top height are often a function of the penetration depth or IR compared with active sensors
- CTT shows less discrepancy than CTP
- Sampling effects could be another factor affecting comparisons (NB use of cost to remove some multi-layer)





FUTURE PLANS

(of particular relevance to fraction and height retrievals)

- More unified aerosol/cloud retrieval scheme
- Improved land surface model
- Move away from simple use of cloud flag, i.e. retrieve all scenes and decide post-hoc if cloud/aerosol using cost of each model + other information (e.g. spatial coherence)
- ATSR forward view in cloud retrieval (PhDs at Oxford and UCL-Stereo retrieval)
- Improved treatment of boundary layer inversions
- Improved treatment of multi-layer systems
 - Recent EUMetSat study
- Synergy between different instruments (AATSR/MERIS, SLSTR/OLCI)- with FUB



Modelling multi layer cloud



- OCA-Optimal Cloud Analysis code for SEVIRI cloud retrievals used to test multi layer cloud model
- Validated with Cloud sat and Calipso data
- R. Siddans et al 2009. EUMETSAT report

OPTIMAL ESTIMATION: APPROACH


- Minimise cost function, J

$$J = (\mathbf{y}_x - \mathbf{y})\mathbf{S}_y^{-1}(\mathbf{y}_x - \mathbf{y})^T + (\mathbf{x} - \mathbf{x}_a)\mathbf{S}_a^{-1}(\mathbf{x} - \mathbf{x}_a)^T$$

- Linearise forward model

$$\mathbf{y}_x = \mathbf{F}\mathbf{x} \quad \delta\mathbf{y}_x = \mathbf{K}\delta\mathbf{x}$$

- Iterative steps weighted between gradient of J and Newtonian descent
 - Stop when change in cost small
- Error estimate on retrieved state



$$J' = \mathbf{K}^T \mathbf{S}_y^{-1}(\mathbf{y}_x - \mathbf{y}) + \mathbf{S}_x^{-1}(\mathbf{x} - \mathbf{x}_a)$$


$$\mathbf{S}_x = \left(\mathbf{S}_a^{-1} + \mathbf{K}^T \mathbf{S}_y^{-1} \mathbf{K} \right)^{-1}$$

Cloud Top Temperature


— ATSR2
— AATSR
— PATMOSX
— ISCCP
— AIRS

— MODIS-STT
— MODIS-CET
— MISR
— POLDER
— CALIPSO

Water CTT



High Ice CTT



- End

